arXiv:cond-mat/9502022v1 6 Feb 1995

Existence of Low-temperature Critical Regime in a One-Dimensional Luttinger Liquid with a Weak Link.

A. M. Tsvelik

February 1, 2008

Department of Physics, University of Oxford, 1 Keble Road, Oxford, OX1 3NP, UK

Abstract

The exact solution of the boundary sine-Gordon model is studied in the region where the scaling dimension of the boundary field $1/2 < \Delta < 1$. It is shown that at $1 > \Delta \geq 2/3$ the infrared fixed fixed point belongs to the universality class of the underscreened anisotrpic Kondo model. The boundary contribution to the specific heat in this region scales as $C \sim T^{2\Delta^{-1}-2}$ at small temperatures.

PACS numbers: 72.10.Fk, 72.15.Qm, 73.20. Dx

The problem of potential scattering in Luttinger liquids has attracted a great deal of attention since Kane and Fisher[1] mapped it onto the Schmid model[2]. The latter model is described by the following action:

$$S = S_0 + M \int \mathrm{d}\tau \cos\left(\beta\phi(0,\tau)/2\right)$$
$$S_0 = \int \mathrm{d}\tau \left\{ \int_0^L \mathrm{d}x \left[\frac{1}{2}(\partial_\tau \phi)^2 + \frac{1}{2}(\partial_x \phi)^2 \right] \right\} \qquad (1)$$

Here the parameter β is determined by interactions in the bulk. It is believed that in the region $\Delta = \beta^2/8\pi < 1$ where the cosine term is relevant the model scales to the strong coupling fixed point described by the effective action

$$S_{\mathrm{eff}} = S_0 + \frac{1}{2} T_B \int \mathrm{d}\tau \phi(0,\tau)^2 \qquad (2)$$

where

$$T_B = M(M/\Lambda)^{\Delta/(1-\Delta)}$$

and Λ is the ultraviolet cut-off. The robustness of the effective action (2) has been proven by the instanton expansion for $\Delta << 1$ [1,3], by the transformation to the equivalent free fermion model ($\Delta = 1/2$)[4,5] and by the exact solution ($\Delta \leq 1/2$) [6,7]. In the entire region described by the effective action (2) the two-point correlation functions on the boundary decay at T = 0 as $1/\tau^2$ ($\tau >> T_B^{-1}$) and the boundary contribution to the specific heat is linear in temperature [8,9]. However, since all these results have been derived only for $\Delta \leq 1/2$, there is still room for something new in the region $1/2 < \Delta < 1$.

In the present Letter we present arguments that in the region $2/3 \leq \Delta < 1$ the model (1) scales to a different critical point where the ϕ^2-term does not appear in the low-energy effective action. In this region the boundary contribution to the specific heat behaves at low temperatures as

$$C_{boundary} \sim T^{2/\Delta - 2}\,(\Delta > 2/3),\ \sim T \ln T\,(\Delta = 2/3) \tag{3}$$

This behaviour is compatible with the presence on the boundary of an irrelevant operator with scaling dimension $1/\Delta > 1$. The latter is the scaling dimension of the dual exponent $\cos\left[2\pi\theta(0)/\beta\right]$ (θ is the field dual to ϕ). This dual exponent is generated by the instanton expansion of the $\cos(\beta\phi/2)$-potential and is always present in the low-energy effective action. However, for $\Delta < 2/3$ its contribution is less important than that of the ϕ^2-term. On the contrary, in the region $2/3 \leq \Delta < 1$ the instanton processes completely screen the parabolic potential. Meanwhile, the operator $\cos(\beta\phi(0)/2)$ remains relevant at high energies throughout the entire region $0 < \Delta < 1$. We suggest that the low energy fixed point in the region $2/3 \leq \Delta < 1$ is described by the effective action

$$S_{\rm eff} = S_0 + \tilde{T}_B \int {\rm d}\tau \cos\left(2\pi\theta(0,\tau)/\beta\right) \tag{4}$$

where $\tilde{T}_B$ is an energy scale related to T_B.

Despite the fact that the boundary sine-Gordon (BSG) model (1) has been solved exactly in the sense that exact S-matrices have been found[6], complexity of the solution in the area $\Delta > 1/2$ has prevented it from beeing studied. At $\Delta = 1/\nu$, $\nu = 2, 3, ...$ the thermodynamic Bethe ansatz (TBA) equations were obtained by Fendley *et al.*[7]:

$$\epsilon_n(v) = \eta\delta_{n,1}\exp(-\pi v/2)$$

$$+s*\ln\left[1+e^{\epsilon_{n-1}(v)}\right]\left[1+e^{\epsilon_{n+1}(v)}\right]+\delta_{n,\nu-2}s*\ln\left[1+e^{\epsilon_\nu(v)}\right]\ (n=1,...\nu-1)$$

$$\epsilon_\nu(v)=s*\ln\left[1+e^{\epsilon_{\nu-2}(v)}\right] \quad (5)$$

$$F_{imp}=-T\int_{-\infty}^{\infty}s[v+\frac{2}{\pi}\ln(T_B/T)]\ln\left[1+e^{\epsilon_{\nu-1}(v)}\right] \quad (6)$$

$$\frac{1}{L}F_{bulk}=-T\int_{-\infty}^{\infty}s[v+\frac{2}{\pi}\ln(\Lambda/T)]\ln\left[1+e^{\epsilon_1(v)}\right] \quad (7)$$

with $\eta=+1$. Here

$$s*f(v)=\int_{-\infty}^{\infty}du\frac{f(u)}{4\cosh[\pi(v-u)/2]}$$

Eqs.(5) are very similar to the equations for the conventional sine-Gordon model. In the latter case there is a duality: TBA equations are invariant under the transformation

$$\Delta\to 1-\Delta=1/\nu \quad (8)$$

except for the free term in the first equation (5), which changes its sign: $\eta=-1$. Therefore we suggest that TBA equations for BSG problem at $\Delta=1-1/\nu$ are given by Eqs.(5, 6) with $\eta=-1$. We have two additional arguments in favour of this proposal. The first one is that the suggested transformation works for the anisotropic spin-1/2 Heisenberg chain with open boundary conditions[8]. It is widely believed that the latter model adequately describes the strong coupling point of BSG model. Indeed, at strong coupling the scattering potential becomes infinite which effectively breaks the chain. The equivalency between the Luttinger liquid of spinless fermions and the spin-1/2 Heisenberg chain is established by the Jordan-Wigner transformation. The second argument is that at $\Delta=3/4$ one can derive TBA equations for BSG model using its equivalency with the 4-channel anisotropic Kondo model in the Toulouse limit[9]. We shall present the latter argument in detail.

The Bethe ansatz equations for an anisotropic k-channel Kondo model are given by

$$[e_k(\eta;u_a)]^N e_{2S}(\eta;u_a-1/g)=\prod_{b=1}^{M}e_2(\eta;u_a-u_b) \quad (9)$$

$$E=\sum_{a=1}^{M}\frac{1}{2\mathrm{i}}\ln e_k(\eta;u_a) \quad (10)$$

$$e_n(\eta;u)=\frac{\sinh[\eta(u-\mathrm{i}n)]}{\sinh[\eta(u+\mathrm{i}n)]} \quad (11)$$

where S is the impurity spin, g is the Kondo coupling constant, η is the anisotropy, N is the length of the system and $M=kN/2+S-S^z$ is the number of up spins.

The universal relationship beetween the quantities g and η and the parameters of the Hamiltonian exists only in the limit of weak anisotropy $\eta << 1$. The difficulty determining the Toulouse limit is resolved if we suggest that this limit corresponds to the maximal value of η at which the IR fixed point of the Kondo model still belongs to the same universality class as at $\eta \to 0$. The periodicity of the trigonometric factors in Eqs.(9) suggests that the Toulouse limit is realized at $\eta = \pi/2(k + 1/\nu)$ ($\nu \to \infty$). This limit has been considered in Refs. 10, 11 and the detailed derivation of TBA equations is given in Ref. 12. It was shown that TBA equations in the Toulouse limit have the following form:

$$\epsilon_n = s * \ln\left(1 + \mathrm{e}^{\epsilon_{n-1}}\right)\left(1 + \mathrm{e}^{\epsilon_{n+1}}\right) + \delta_{n,k-2} s * \ln\left(1 + \mathrm{e}^{\epsilon_k}\right),\ n = 1, ...k-1 \qquad (12)$$

$$\epsilon_k = s * \ln\left(1 + \mathrm{e}^{\epsilon_{k-2}}\right) - 2\exp(-\pi v/2) \qquad (13)$$

$$F_{bulk} = -NT^2 \int_{-\infty}^{\infty} \mathrm{d}v \mathrm{e}^{-\pi v/2} \ln\left(1 + \mathrm{e}^{\epsilon_k}\right) \qquad (14)$$

$$F_{boundary} = -T \int_{-\infty}^{\infty} \mathrm{d}v s(v + \frac{2}{\pi} \ln T_B/T) \ln\left(1 + \mathrm{e}^{\epsilon_{2S}}\right) \qquad (15)$$

where the temperature T_B is defined as

$$T_B = \lim_{\nu,\Lambda\to\infty} \frac{\pi\Lambda}{2\nu} \exp(-\pi/2g)$$

Executing this limit one has to be careful to keep the energy scale T_B finite. Substituting into these equations $S = 1/2$ and $k = 4$ we reproduce the suggested equations for the BSG model with $\Delta = 3/4$ (i.e. $\nu = 4$, $\eta = -1$). The fact that this equivalency holds only for $k = 4$ is not suprising. The conformal charge of the bulk degrees of freedom in the k-channel Kondo model is equal to[13,14]

$$C = \frac{3k}{k+2} \qquad (16)$$

At $k = 4$ it is equal to 2 which corresponds to two bosonic modes interacting with the impurity. In the Toulouse limit one of these modes decouples from the impurity and the effective conformal charge becomes 1.

Now we shall calculate the IR and UV asymptotics of the boundary free energy for the BSG model with $\Delta = 1 - 1/\nu$. Analytical solutions are available for asymptotics of $\epsilon_n(v)$ at $v \to \pm\infty$ (see, for example, Ref.(15)). At large temperatures $T >> T_B$ the free energy is determined by the asymptotics at $v \to +\infty$ where

$$\left(1 + \mathrm{e}^{\epsilon_{\nu-1}}\right) = \nu \qquad (17)$$

so that we have

$$F_{boundary} \to -\frac{T}{2}\ln\nu \tag{18}$$

At small temperatures the leading contribution comes from the region $v \to 0$ where ϵ_n ($\nu \neq 1$) are again almost constant and $\exp(\epsilon_1)$ is small. Then the corrections can be determined from the expansion in $\exp(\epsilon_1)$:

$$g_n(v) \equiv \ln\left(1+\mathrm{e}^{\epsilon_n(v)}\right) = g_n^{(0)} + g_n^{(1)}(v) + ...$$

$$g_n^{(0)} = 2\ln n \ (n = 2, 3, ...\nu - 2),\ g_{\nu-1}^{(0)} = g_\nu^{(0)} \ln(\nu - 1)$$

$$g_n^{(1)}(v) = \frac{1}{2n}[(n+1)a_n * g_1(v) - (n-1)a_{n+2} * g_1(v)] \tag{19}$$

$$a_n(\omega) = \frac{\sinh(\nu - n)\omega}{\sinh(\nu - 1)\omega} \tag{20}$$

Using these expressions we get the following expansion for the free energy:

$$F_{boundary} \to -\frac{T}{2}\ln(\nu-1) - T(\nu-1)\int_0^\infty \mathrm{d}v f(v)\ln\left(1+\mathrm{e}^{\epsilon_1(v)}\right) \tag{21}$$

where

$$f(\omega) = \frac{\tanh\omega}{\sinh(\nu-1)\omega}$$

The first term in Eq.(21) gives the finite entropy of the ground state $S(0) = -\frac{1}{2}\ln(\Delta^{-1}-1)$. A careful analysis shows that this entropy disappears at $\Delta \leq 1/2$. The ratio of the partition functions in the ultraviolet and the infrared limits is

$$Z_{UV}/Z_{IR} = \Delta^{-1/2} \tag{22}$$

This result reproduces the expression obtained for $\Delta < 1/2$ in Ref. 7. The second term in Eq.(21) gives $\delta F_{imp} \sim -T^{1+2/(\nu-1)}$ which leads to Eq.(3).

The author expresses his gratitude to P. Coleman, M. Evans, E. Fradkin, L. Ioffe and P. de Sa for the valuable discussions and the interest to the work.

References

[1] C. L. Kane and M. P. A. Fisher, Phys. Rev. Lett. **68**, 1220 (1992); Phys. Rev. **B46**, 15233 (1992).

[2] A. Schmid, Phys. Rev. Lett. **51**, 1506 (1983).

[3] M. P. A. Fisher and W. Zwerger, Phys. Rev. B**32**, 6190 (1985).

[4] F. Guinea, Phys. Rev. B**32**, 7518 (1985).

[5] K. A. Matveev, Phys. Rev. B.**51**, 1743 (1995).

[6] S. Ghoshal and A. B. Zamolodchikov, Int. J. Mod. Phys. **A9**, 3841 (1994).

[7] P. Fendley, H. Saleur and N. P. Warner, to appear in Nucl. Phys. B, sissa preprint hep-th/9406125.

[8] P. de Sa and A. M. Tsvelik, unpublished.

[9] Fabrizio and A. Gogolin, to appear in Phys. Rev. B.

[10] P. B. Wiegmann, Pis'ma Zh. Eksp. Teor. Fiz. **41**, 79 (1984) [JETP Lett. **41**, 95 (1984)]; Phys. Lett. B**141**, 217 (1984).

[11] A. N. Kirillov and N. Yu. Reshetikhin, Zapiski Nauch. Seminara LOMI **145**, 109; **146**, 47 (1985).

[12] A. M. Tsvelik, Zh. Eksp. Teor. Fiz. **93**, 385 (1987) [JETP **66**, 221 (1987)].

[13] A. M. Tsvelik, J. Phys. C**18**, 159 (1985).

[14] I. Affleck and A. W. W. Ludwig, Nucl. Phys. B**360**, 641 (1991).

[15] M. Takahashi, M. Suzuki, Prog. Theor. Phys. **48**, 2187 (1972).